



ST LEONARDS DERMATOLOGY & LASER

WOUND CARE BIOPSY

Do not wet the wound for 24 hours.

After 24 hours you may wet the wound. If the wound gets too saturated then pat it dry with a clean tissue. Leave the dressing on for up to 4 days or until you return for removal of stitches.

Do the following if you wish to change the dressing:

- 1) Clean wound with one teaspoon of salt added to one cup of water using a piece of gauze or clean cotton tip. Then pat dry with with a clean tissue.
- 3) Apply a small amount of Vaseline to the wound/suture site.
- 4) Place a piece of Hypafix dressing (supplied) along the suture/wound.
- 5) It is not necessary to do this daily, as it is best if the wound is not disturbed too much.
- 6) Signs of infection include pain, swelling, redness or discharge present in the area. See below

The area may drain clear fluid, and you may notice some white material in the base of the wound and a thin rim of redness around the wound as it heals. The wound may also develop a crust or scab. These developments are normal.

If the wound becomes very tender or swollen with increasing redness, or if pus drains from the wound, this may be a sign of infection. If this occurs, please call us on **(02) 99669667** during business hours.

If wound continues to bleed, please apply firm pressure with a clean cloth or gauze and elevate the area if possible for 20-30minutes. If the bleeding does not stop then please contact us on the numbers above.